

MSC2010: 53C55, 53B35

© *F. H. Al-Hussaini, H. M. Abood*

## QUAISI INVARIANT CONHARMONIC TENSOR OF SPECIAL CLASSES OF LOCALLY CONFORMAL ALMOST COSYMPLECTIC MANIFOLD

The authors classified a locally conformal almost cosymplectic manifold ( $\mathcal{LCA}C_S$ -manifold) according to the conharmonic curvature tensor. In particular, they have determined the necessary conditions for a conharmonic curvature tensor on the  $\mathcal{LCA}C_S$ -manifold of classes  $CT_i, i = 1, 2, 3$  to be  $\Phi$ -quasi invariant. Moreover, it has been proved that any  $\mathcal{LCA}C_S$ -manifold of the class  $CT_1$  is conharmonically  $\Phi$ -paracontact.

*Keywords:* locally conformal almost cosymplectic manifold, conharmonic curvature tensor,  $\Phi$ -quasi invariant, conharmonically  $\Phi$ -paracontact.

DOI: [10.35634/vm200201](https://doi.org/10.35634/vm200201)

### Introduction

The first classification of an almost contact metric structure has been introduced by Chinea and Marrero [9]. Like an almost Hermitian structure, another principle of classification of an almost contact metric structure for differential-geometric invariants of the second order depends on the properties of the Riemannian curvature tensor  $R$ . Volkova introduced analogues of these classes in contact geometry  $CR_1, CR_2$  and  $CR_3$  [19].

Our study focuses on the conharmonic tensor of the locally conformal almost cosymplectic manifold of three special classes  $CT_i, i = 1, 2, 3$  which are related with the classes  $CR_1, CR_2, CR_3$ .

A harmonic function is a function whose Laplacian vanishes. It is known that the conformal transformation on the Riemannian manifold preserves the angle between two vectors. Generally, a harmonic function is not invariant. The condition to remain such a function invariant has been studied by Ishi [13]. Specifically, he introduced a conharmonic transformation that preserves the harmonicity of a certain function. On the other hand, Ghosh et al. [11] studied the  $N(k)$ -contact metric manifolds satisfying curvature conditions on the conharmonic tensor. Dwivedi and Kim [10] obtained certain necessary and sufficient conditions for the  $K$ -contact and Sasakian manifolds to be quasi conharmonically flat,  $\xi$ -conharmonically flat and  $\Phi$ -conharmonically flat. Furthermore, Asghari and Taleshian [4] studied the conharmonic curvature tensor on the Kenmotsu manifold. Chanyal and Uperti [8] proved that  $\Phi$ -conharmonically flat  $(k, \mu)$ -contact manifolds are  $\eta$ -Einstein manifolds. Taleshian et al. [18] considered  $LP$ -Sasakian manifolds admitting a conharmonic curvature tensor. Abood and Al-Hussaini [3] studied the geometrical properties of the conharmonic tensor of a locally conformal almost cosymplectic manifold. In particular, the authors established the necessary and sufficient conditions for the conharmonic tensor to be flat, the aforementioned manifold to be normal and an  $\eta$ -Einstein manifold.

### § 1. Preliminaries

The main purpose of this section is to construct an almost contact metric structure and a locally conformal almost cosymplectic manifold in the adjoined  $G$ -structure space.

**Definition 1** (see [5]). Let  $M$  be  $2n + 1$  dimensional smooth manifold,  $\eta$  be differential 1-form called a *contact form*,  $\xi$  be a vector field called a *characteristic*,  $\Phi$  be an endomorphism of the module of the vector fields  $X(M)$  called a *structure endomorphism*. The family of tensors  $\{\eta, \xi, \Phi\}$  is called an *almost contact structure* if the following conditions hold

1.  $\eta(\xi) = 1$ ;
2.  $\Phi(\xi) = 0$ ;
3.  $\eta \circ \Phi = 0$ ;
4.  $\Phi^2 = -id + \eta \otimes \xi$ .

In addition, if there is a Riemannian metric  $g = \langle \cdot, \cdot \rangle$  on  $M$  such that

$$\langle \Phi X, \Phi Y \rangle = \langle X, Y \rangle - \eta(X)\eta(Y), \quad X, Y \in X(M),$$

then the family of tensors  $\{\eta, \xi, \Phi, g\}$  is called an almost contact metric structure. In this case, the mentioned manifold  $M$  endowed with this structure is called an almost contact metric manifold.

**Definition 2** (see [16]). At each point  $p \in M^{2n+1}$ , there is a frame in a complexification of a tangent space  $T_p^c(M)$  of the form  $(p, \varepsilon_0, \varepsilon_1, \dots, \varepsilon_n, \varepsilon_{\hat{1}}, \dots, \varepsilon_{\hat{n}})$ , where  $\varepsilon_a = \sqrt{2}\pi(e_a)$ ,  $\varepsilon_{\hat{a}} = \sqrt{2}\bar{\pi}(e_a)$ ,  $\hat{a} = a + n$ ,  $\varepsilon_0 = \xi_p$ ,  $\pi = -\frac{1}{2}(\Phi^2 + \sqrt{-1}\Phi)$  and  $\bar{\pi} = \frac{1}{2}(-\Phi^2 + \sqrt{-1}\Phi)$ . The frame  $(p, \varepsilon_0, \varepsilon_1, \dots, \varepsilon_n, \varepsilon_{\hat{1}}, \dots, \varepsilon_{\hat{n}})$  is called an  $A$ -frame.

A set of such frames defines  $G$ -structure on  $M$  with a structure group  $1 \times U(n)$ . This structure is called a  $G$ -adjoined structure.

**Lemma 1** (see [15]). The component matrices of the tensors  $\Phi_p$  and  $g_p$  in  $A$ -frame have the following forms respectively:

$$(\Phi_j^i) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \sqrt{-1}I_n & 0 \\ 0 & 0 & -\sqrt{-1}I_n \end{pmatrix}, \quad (g_{ij}) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & I_n \\ 0 & I_n & 0 \end{pmatrix},$$

where  $I_n$  is the identity matrix of order  $n$ .

**Definition 3** (see [5]). A skew-symmetric tensor  $\Omega(X, Y) = g(X, \Phi Y)$  is called a fundamental form of the  $\mathcal{AC}$ -structure.

**Definition 4** (see [12]). An almost contact metric structure  $S = (\eta, \xi, \Phi, g)$  is called an almost cosymplectic structure ( $\mathcal{AC}_f$ -structure) if

1.  $d\eta = 0$ ;
2.  $d\Omega = 0$ .

**Definition 5** (see [17]). A conformal transformation of an  $\mathcal{AC}$ -structure  $S = (\eta, \xi, \Phi, g)$  on a manifold  $M$  is a mapping from  $S$  to an  $\mathcal{AC}$ -structure  $\tilde{S} = (\tilde{\eta}, \tilde{\xi}, \tilde{\Phi}, \tilde{g})$  such that

$$\tilde{\eta} = e^{-\sigma}\eta, \quad \tilde{\xi} = e^{\sigma}\xi, \quad \tilde{\Phi} = \Phi, \quad \tilde{g} = e^{-2\sigma}g,$$

where  $\sigma$  is the determining function of the conformal transformation. If  $\sigma = \text{const}$ , then the conformal transformation is said to be trivial.

**Definition 6** (see [17]). An  $\mathcal{AC}$ -structure  $S$  on a manifold  $M$  is said to be a locally conformal almost cosymplectic ( $\mathcal{LCA}_S$ -structure) if the restriction of  $S$  on a some neighborhood  $U$  of an arbitrary point  $p \in M$ , admits a conformal transformation of an almost cosymplectic structure. This transformation is called a locally conformal. A manifold  $M$  equipped with a  $\mathcal{LCA}_S$ -structure is called a  $\mathcal{LCA}_S$ -manifold.

**Lemma 2** (see [14]). *In the adjoined  $G$ -structure space, the structure equations of  $\mathcal{LCA}_S$ -manifold have the following forms:*

1.  $d\omega^a = -\omega_b^a \wedge \omega^b + B_c^{ab}\omega^c \wedge \omega_b + B^{abc}\omega_b \wedge \omega_c + B_b^a\omega \wedge \omega^b + B^{ab}\omega \wedge \omega_b;$
2.  $d\omega_a = \omega_a^b \wedge \omega_b + B_{ab}^c\omega_c \wedge \omega^b + B_{abc}\omega^b \wedge \omega^c + B_a^b\omega \wedge \omega_b + B_{ab}\omega \wedge \omega^b;$
3.  $d\omega = C_b\omega \wedge \omega^b + C^b\omega \wedge \omega_b;$

where  $A_{[bd]}^{ac} = -2\delta_{[b}^{[c}\sigma_{d]}^a] + 2\sigma^{[a}\delta_b^{e]}\sigma_{[c}\delta_{d]}^e + \frac{1}{2}B^{dca}B_{ebd}$ , Here  $B^{abc}$ ,  $B_{abc}$ ;  $B^{ab}$ ,  $B_{ab}$ ;  $B_b^a$ ,  $B_a^b$ ;  $C^{ab}$ ,  $C_{ab}$ ;  $C^b$ ,  $C_b$ ;  $A_b^{acd}$ ,  $A_{acd}^b$ ;  $A_{bd}^{ac}$ ;  $A_b^{ac0}$ ,  $A_{ac0}^b$ ;  $B^{abci}$ ,  $B_{abci}$ ;  $D^{abi}$ ,  $D_{abi}$  and  $\sigma_{ij}$  are smooth functions in the adjoined  $G$ -structure space. The tensors  $B^{abc}$ ;  $B^{ab}$  are called the second and third structure tensors respectively.

**Definition 7** (see [7]). A Ricci tensor is a tensor of type  $(2, 0)$  which is a contracting of the Riemannian curvature tensor and defined by

$$r_{ij} = -R_{ijk}^k.$$

**Lemma 3** (see [1]). *In the adjoined  $G$ -structure space, the components of the Ricci tensor of  $\mathcal{LCA}_S$ -manifold are given by the following forms:*

1.  $r_{ab} = 2(-2A_{(ab)c}^c - 4(\sigma^{[c}\delta_{[b}^{h]}B_{c]ha} + \sigma^{[c}\delta_{[a}^{h]}B_{c]hb}) + \sigma_0 B_{a[c}\delta_{b]}^c + \sigma_0 B_{b[c}\delta_{a]}^c + 2\sigma_0 B_{ab} - D_{ab0} - \sigma_{ab} - \sigma_a\sigma_b + 2B_{bah}\sigma^h;$
2.  $r_{\dot{a}\dot{b}} = -4(\delta_{[b}^{[a}\sigma_{c]}^c] - \sigma_{[c}\delta_{[b}^{h]}\sigma^{[h}\delta_c^a] - \frac{1}{2}\sigma^{[a}\delta_b^{h]}\sigma_h + B^{hca}B_{hcb} + B^{bch}B_{cha}) + (B^{cb}B_{ac} - B_{hb}B^{ah}) + A_{ac}^{cb} - \delta_b^a\sigma_{00} - 2n\sigma_0^2 - \sigma_b^a - \sigma^a\sigma_b;$
3.  $r_{a0} = -A_{ac0}^c - \sigma^c B_{ac} + n\sigma_0\sigma_a + 2(\sigma_{0[c}\delta_{a]}^c + B^{cb}B_{bca} - 2\sigma^{[c}\delta_{[c}^{h]}B_{a]h});$
4.  $r_{oo} = -2n(\sigma_{00} + \sigma_0^2) - 2B_{hc}B^{ch} - 2(\sigma_c^c + \sigma^c\sigma_c) + 4\sigma^{[c}\delta_c^{h]}\sigma_h.$

The remaining components are conjugate to the above-mentioned components.

**Definition 8** (see [2]). A  $\mathcal{LCA}_S$ -manifold is said to have  $\Phi$ -invariant Ricci tensor, if  $\Phi \circ r = r \circ \Phi$ .

**Lemma 4** (see [2]). *A  $\mathcal{LCA}_S$ -manifold has  $\Phi$ -invariant Ricci tensor if and only if, in the adjoined  $G$ -structure space, the following condition*

$$r_b^{\dot{a}} = r_{ab} = r_0^{\dot{a}} = r_{a0} = 0$$

holds.

**Definition 9** (see [6]). A pseudo-Riemannian manifold  $M$  is known as an  $\eta$ -Einstein of type  $(\alpha, \beta)$  if its Ricci tensor satisfies the following condition:

$$r = \alpha g + \beta \eta \otimes \eta,$$

where  $\alpha$  and  $\beta$  are suitable smooth functions. If  $\beta = 0$ , then  $M$  is referred to as an Einstein manifold.

**Definition 10** (see [12]). Let  $M$  be an  $\mathcal{AC}$ -manifold of dimension  $2n + 1$ . A tensor  $T$  of type  $(4, 0)$  which is invariant under conharmonic transformation and defined by the form

$$T_{ijkl} = R_{ijkl} - \frac{1}{2n-1}(r_{il}g_{jk} - r_{jl}g_{ik} + r_{jk}g_{il} - r_{ik}g_{jl})$$

is called a conharmonic curvature tensor.

In the work [1], we have calculated all possible components of the mentioned tensor which are listed in the next lemma.

**Lemma 5.** *In the adjoined  $G$ -structure space, the components of the conharmonic curvature tensor of  $\mathcal{LCA}_S$ -manifold are given by the following forms:*

1.  $T_{abcd} = 2(2B_{[c|ab|d]} - 2\sigma_{[a}B_{b]cd} + B_{a[c}B_{d]b});$
2.  $T_{\hat{a}bcd} = 2(A_{bcd}^a + 4\sigma^{[a}\delta_{[c}^h]B_{d]hb} - \sigma_0 B_{b[d}\delta_{c]}^a) - \frac{1}{2n-1}(r_{bc}\delta_d^a - r_{bd}\delta_c^a);$
3.  $T_{\hat{a}bc\hat{d}} = A_{bc}^{ad} + 4\sigma^{[a}\delta_c^h]\sigma_{[h}\delta_b^d] - 4B^{dah}B_{chb} + B^{ad}B_{bc} - \delta_c^a\delta_b^d\sigma_0^2 + \frac{1}{2n-1}(r_b^d\delta_c^a + r_c^a\delta_b^d);$
4.  $T_{\hat{a}\hat{b}cd} = 2(2\delta_{[c}^{[b}\sigma_{d]}^a] + 2B^{hab}B_{hdc} - \delta_{[c}^a\delta_{d]}^b\sigma_0^2) - \frac{4}{2n-1}(r_{[a}^{[d}\delta_{b]}^c]);$
5.  $T_{\hat{a}0cd} = 2(\sigma_{0[c}\delta_{d]}^a + B^{ab}B_{bcd} - 2\sigma^{[a}\delta_{[c}^h]B_{d]h}) + \frac{1}{2n-1}(r_{0d}\delta_c^a - r_{0c}\delta_d^a);$
6.  $T_{\hat{a}\hat{b}c0} = A_b^{ac0} + \sigma_b B^{ac} - \delta_b^c\sigma_0\sigma^a - \frac{1}{2n-1}(r_0^a\delta_b^c);$
7.  $T_{abc0} = 2B_{cab0} + 2B_{cab}\sigma_0;$
8.  $T_{\hat{a}0b0} = -\delta_b^a\sigma_{00} - \delta_b^a\sigma_0^2 - B_{cb}B^{ac} - \sigma_b^a - \sigma^a\sigma_b + 2\sigma^{[a}\delta_b^c]\sigma_c + \frac{1}{2n-1}(r_{00}\delta_b^a + r_b^a);$
9.  $T_{\hat{a}0\hat{b}0} = 2\sigma_0 B^{ab} - D^{ab0} - \sigma^{ab} - \sigma^a\sigma^b + 2B^{bac}\sigma_c + \frac{1}{2n-1}(r_{\hat{a}\hat{b}}).$

*The remaining components are conjugate to the above-mentioned components or can be obtained by the property of symmetry for  $T$  or identically equal to zero.*

The next lemma gives analogues to the Gray's identities in the adjoined  $G$ -structure space.

**Lemma 6** (see [19]). *An  $\mathcal{AC}$ -manifold is called a manifold of class*

1.  $CR_1$  if and only if,  $R_{abcd} = R_{\hat{a}bcd} = R_{\hat{a}\hat{b}cd} = 0;$
2.  $CR_2$  if and only if,  $R_{abcd} = R_{\hat{a}bcd} = 0;$
3.  $CR_3$  if and only if,  $R_{\hat{a}bcd} = 0.$

## § 2. The main results

In this section, we introduce analogues to the Gray's identities for the conharmonic curvature tensor of a  $\mathcal{LCA}_S$ -manifold.

**Definition 11.** In the adjoined  $G$ -structure space, a  $\mathcal{LCA}_S$ -manifold is called a manifold of class

1.  $CT_1$  if and only if,  $T_{abcd} = T_{\hat{a}bcd} = T_{\hat{a}\hat{b}cd} = 0;$
2.  $CT_2$  if and only if,  $T_{abcd} = T_{\hat{a}bcd} = 0;$

3.  $CT_3$  if and only if,  $T_{abcd} = 0$ .

**Definition 12.** The conharmonic tensor is called  $\Phi$ -quasi invariant , if

$$\begin{aligned} \langle T(\Phi X, \Phi Y)\Phi X, \Phi Y \rangle - \langle T(\Phi^2 X, \Phi^2 Y)\Phi^2 X, \Phi^2 Y \rangle &= \langle T(\Phi^2 X, \Phi Y)\Phi^2 X, \Phi Y \rangle \\ &- \langle T(\Phi X, \Phi^2 Y)\Phi X, \Phi^2 Y \rangle \end{aligned}$$

holds for each  $X, Y, Z \in X(M)$ .

**Definition 13.** A  $\mathcal{LCA}_S$ -manifold is called a conharmonically  $\Phi$ -paracontact, if its conharmonic tensor satisfies the identity

$$\begin{aligned} \langle T(\Phi^2 X, \Phi^2 Y)\Phi^2 Z, \Phi^2 W \rangle &= \langle T(\Phi^2 X, \Phi^2 Y)\Phi Z, \Phi W \rangle - \langle T(\Phi X, \Phi^2 Y)\Phi Z, \Phi^2 W \rangle \\ &- \langle T(\Phi X, \Phi^2 Y)\Phi^2 Z, \Phi W \rangle, \end{aligned}$$

where  $X, Y, \in X(M)$ .

**Theorem 1.** A conharmonic tensor on  $\mathcal{LCA}_S$ -manifold of class  $CT_i$ ,  $i = 1, 2, 3$ , is  $\Phi$ -quasi invariant.

**P r o o f.** Consider the component  $T_{\bar{a}bab} = 0$ . Write this equation in  $A$ -frame, we have

$$\langle T(\varepsilon_{\bar{a}}, \varepsilon_b)\varepsilon_a, \varepsilon_b \rangle = 0.$$

Then we get

$$\langle T(\bar{\sigma}X, \sigma Y)\sigma X, \sigma Y \rangle = 0, \quad X, Y \in X(M)$$

We compensate for the value of the projections  $\sigma$  and  $\bar{\sigma}$  and, using the linear property of the tensor, we can rewrite the previous expression in the form

$$\begin{aligned} &\{ \langle T(X, Y)X, Y \rangle + \langle T(\Phi X, \Phi Y)X, Y \rangle + \langle T(\Phi X, Y)\Phi X, Y \rangle + \langle T(\Phi X, Y)X, \Phi Y \rangle - \\ &\langle T(X, \Phi Y)\Phi X, Y \rangle - \langle T(X, Y)\Phi X, \Phi Y \rangle - \langle T(X, \Phi Y)X, \Phi Y \rangle - \langle T(\Phi X, \Phi Y)\Phi X, \Phi Y \rangle \} + \\ &i \{ \langle T(X, \Phi Y)\Phi X, \Phi Y \rangle - \langle T(X, \Phi Y)X, Y \rangle - \langle T(X, Y)\Phi X, Y \rangle - \langle T(X, Y)X, \Phi Y \rangle + \\ &\langle T(\Phi X, Y)X, Y \rangle - \langle T(\Phi X, \Phi Y)\Phi X, Y \rangle - \langle T(\Phi X, \Phi Y)X, \Phi Y \rangle - \langle T(\Phi X, Y)\Phi X, \Phi Y \rangle \} = 0. \end{aligned}$$

Note that the real part equals to zero, and regarding the symmetrical properties of the coharmonic curvature tensor, we have

$$\langle T(X, Y)X, Y \rangle + \langle T(\Phi X, Y)\Phi X, Y \rangle - \langle T(X, \Phi Y)X, \Phi Y \rangle - \langle T(\Phi X, \Phi Y)\Phi X, \Phi Y \rangle = 0. \quad (1)$$

Then by substituting  $X \rightarrow -\Phi^2 X; Y \rightarrow -\Phi^2 Y$  in equation (1) and using the fact  $\Phi^3(X) = -\Phi(X)$ , we obtain the assertion of the theorem.  $\square$

**Theorem 2.** A conharmonic curvature tensor on  $\mathcal{LCA}_S$ -manifold of class  $CR_3$  with  $\Phi$ -invariant Ricci tensor is  $\Phi$ -quasi invariant.

**P r o o f.** Since the  $\mathcal{LCA}_S$ -manifold  $M$  has  $\Phi$ -invariant Ricci tensor, then  $CT_3$  and  $CR_3$  are coincide. Regarding the Theorem 1, we get that the conharmonic curvature tensor is  $\Phi$ -quasi invariant.  $\square$

**Theorem 3.** Any  $\mathcal{LCA}_S$ -manifold  $M$  of class  $CT_1$  is conharmonically  $\Phi$ -paracontact.

**P r o o f.** On the  $G$ -adjoined structure space, we have  $T_{\hat{a}bcd} = 0$ . In  $A$ -frame, we get

$$\langle T(\varepsilon_{\hat{a}}, \varepsilon_{\hat{b}})\varepsilon_c, \varepsilon_d \rangle = 0.$$

Consequently, we have

$$\langle T(\bar{\sigma}X, \sigma Y)\sigma X, \sigma Y \rangle = 0, \quad X, Y, Z, W \in X(M).$$

We make up for the value of the projections  $\sigma$  and  $\bar{\sigma}$  and, using the linear property of the tensor, we can rewrite this expression in the form

$$\langle T(X, Y)Z, W \rangle - \langle T(\Phi X, \Phi Y)Z, W \rangle + \langle T(X, \Phi Y)\Phi Z, W \rangle + \langle T(X, \Phi Y)Z, \Phi W \rangle - \langle T(X, Y)\Phi Z, \Phi W \rangle + \langle T(\Phi X, Y)\Phi Z, W \rangle + \langle T(\Phi X, Y)Z, \Phi W \rangle + \langle T(\Phi X, \Phi Y)\Phi Z, \Phi W \rangle = 0. \quad (2)$$

According to the Definition 11, we have  $T_{abcd} = 0$ . Using the same technique, consequently we obtain

$$\langle T(X, Y)Z, W \rangle + \langle T(\Phi X, \Phi Y)Z, W \rangle - \langle T(X, \Phi Y)\Phi Z, W \rangle - \langle T(X, \Phi Y)Z, \Phi W \rangle - \langle T(X, Y)\Phi Z, \Phi W \rangle + \langle T(\Phi X, Y)\Phi Z, W \rangle + \langle T(\Phi X, Y)Z, \Phi W \rangle - \langle T(\Phi X, \Phi Y)\Phi Z, \Phi W \rangle = 0. \quad (3)$$

From the combination of the equations (2) and (3), it follows that

$$\langle T(X, Y)Z, W \rangle = \langle T(X, Y)\Phi Z, \Phi W \rangle - \langle T(\Phi X, Y)\Phi Z, W \rangle - \langle T(\Phi X, Y)Z, \Phi W \rangle. \quad (4)$$

Then by substituting  $X \rightarrow -\Phi^2 X; Y \rightarrow -\Phi^2 Y; Z \rightarrow -\Phi^2 Z, W \rightarrow -\Phi^2 W$  in the equation (4) and using the fact  $\Phi^3(X) = -\Phi(X)$ , we get the result.  $\square$

**Theorem 4.** *Let  $M$  be  $\mathcal{LCA}_S$ -manifold of class  $CR_2$ , then the first structure tensor is parallel in the first canonical connection.*

**P r o o f.** Suppose that  $M$  is  $\mathcal{LCA}_S$ -manifold of the class  $CR_2$ . So we have

$$2((B_{cabd} - B_{dabc}) - 2\sigma_{[a}B_{b]cd} + B_{a[c}B_{d]b}) = 0. \quad (5)$$

Symmetrizing (5) by the indices  $(c, d)$  and then by the indices  $(c, b)$ , we get

$$B_{cabd} = 0.$$

Regarding the fundamental theorem of tensor analysis, we have

$$\nabla B_{abc} = dB_{abc} + B_{dbc}\omega_a^d + B_{adc}\omega_b^d + B_{abd}\omega_c^d = B_{abcd}\omega^d.$$

So, we get

$$\nabla B_{abc} = B_{abcd}\omega^d.$$

It follows that

$$\nabla B_{abc} = 0.$$

Therefore, the tensor  $B_{abc}$  is parallel.  $\square$

**Theorem 5.** *Let  $M$  be  $\mathcal{LCA}_S$ -manifold of class  $CR_1$ , then its second and third structure tensors identically vanish.*

**P r o o f.** Let  $M$  be  $\mathcal{LCA}_S$ -manifold of the class  $CR_1$ . According to the component of Riemannian curvature tensor, we have

$$2(2B[c|ab|d] - 2\sigma_{[a}B_{b]cd}B_{a[c}B_{d]b}) = 0. \tag{6}$$

Symmetrizing and then antisymmetrizing (6) by the indices  $(a, b)$ , we deduce

$$B_{ac}B_{db} - B_{ad}B_{cb} = 0. \tag{7}$$

Antisymmetrizing (7) by the indices  $(a, d)$ , we get

$$B_{ac}B_{db} = 0. \tag{8}$$

Contracting (8) by the indices  $(a, d)$  and then by  $(c, b)$ , we get  $B_{ac}^2 = 0$ , then

$$B_{ac} = 0.$$

From the condition  $R_{\hat{a}\hat{b}\hat{c}\hat{d}} = 0$ , we obtain

$$B^{abc} = 0.$$

□

In the rest of this section, we establish a relation between the classes  $CT_1$  and  $CR_1$ .

**Theorem 6.** *Suppose that  $M$  is  $\mathcal{LCA}_S$ -manifold of class  $CT_1$  with  $\Phi$ -invariant Ricci tensor; then  $M$  is a manifold of class  $CR_1$  if and only if  $M$  has flat Ricci tensor.*

**P r o o f.** Suppose that  $M$  has flat Ricci tensor. Since  $M$  is a  $\mathcal{LCA}_S$ -manifold of class  $CT_1$ , then according to the Definition 11, we have

$$T_{abcd} = T_{\hat{a}\hat{b}\hat{c}\hat{d}} = T_{\hat{a}\hat{b}\hat{c}\hat{d}} = 0.$$

The condition  $T_{abcd} = 0$ , implies that  $R_{abcd} = 0$ ; while the condition  $T_{\hat{a}\hat{b}\hat{c}\hat{d}} = 0$  with  $\Phi$ -invariant Ricci tensor, implies that  $R_{\hat{a}\hat{b}\hat{c}\hat{d}} = 0$ . In addition,  $T_{\hat{a}\hat{b}\hat{c}\hat{d}} = 0$  with flat Ricci tensor, indicates  $R_{\hat{a}\hat{b}\hat{c}\hat{d}} = 0$ . Then  $M$  is a manifold of class  $CR_1$ . Conversely, according to the condition  $T_{\hat{a}\hat{b}\hat{c}\hat{d}} = 0$ , we have

$$2(2\delta_{[c}^{[b}\sigma_{d]}^{a]} + 2B^{hab}B_{hdc} - \delta_{[c}^a\delta_{d]}^b\sigma^2) - \frac{4}{2n-1}(r_{[c}^{[a}\delta_{d]}^{b]}) = 0.$$

Since  $M$  is a manifold of class  $CR_1$ , we get

$$-\frac{4}{2n-1}(r_{[c}^{[a}\delta_{d]}^{b]}) = 0. \tag{9}$$

Taking the contraction operation for (9) by the indices  $(b, d)$ , we deduce

$$r_c^a\delta_b^b - r_b^a\delta_c^b - r_c^b\delta_b^a + r_b^b\delta_c^a = 0.$$

Consequently, we obtain

$$(n-2)r_c^a + r_b^b\delta_c^a = 0. \tag{10}$$

Taking the symmetrization and antisymmetrization operations for (10) by the indices  $(b, a)$ , we conclude

$$r_c^a = 0. \tag{11}$$

Making use of the Theorems 3 and 5, we have

$$-4(\delta_{[b}^a \sigma_{c]}^c] - \sigma_{[c} \delta_{h]}^c \sigma^{[h} \delta_c^a] - \frac{1}{2} \sigma^{[a} \delta_b^h] \sigma_h) + A_{ac}^{cb} - \delta_b^a \sigma_{00} - 2n \delta_b^a \sigma_0^2 - \sigma_b^a - \sigma^a \sigma_b = 0. \quad (12)$$

Symmetrizing (12) by the indices  $(c, h)$ , we obtain

$$-\sigma_{[c} \delta_{h]}^c \sigma^{[h} \delta_c^a] = 0. \quad (13)$$

Contracting (13) by the indices  $(a, c)$ , we get

$$\sigma_a \delta_h^a \sigma^h \delta_a^a - \sigma_a \delta_h^a \sigma^a \delta_a^h - \sigma_h \delta_a^a \sigma^h \delta_a^a + \sigma_h \delta_a^a \sigma^a \delta_a^h = 0.$$

Hence

$$-n(n-1)(\sigma_h \sigma^h) = 0.$$

Or equivalently,

$$\sigma_h \bar{\sigma}_h = 0 \Leftrightarrow \sum_h |\sigma_h|^2 = 0 \Leftrightarrow \sigma_h = 0. \quad (14)$$

Once again, by using the equation (12), we have

$$A_{ac}^{cb} - \delta_b^a \sigma_{00} - 2n \delta_b^a \sigma_0^2 = 0. \quad (15)$$

By virtue of the Theorems 2, 5 and equation (14), the tensors  $A_{[bd]}^{ac} = A_{bd}^{[ac]} = 0$  become symmetric regarding the lower and upper indices. Antisymmetrizing (15) by the indices  $(c, b)$ , we obtain

$$-\delta_b^a \sigma_{00} - 2n \delta_b^a \sigma_0^2 = 0. \quad (16)$$

Contracting (16) by the indices  $(a, b)$ , consequently, we get

$$r_{00} = 0. \quad (17)$$

According to the equations (11), (17) and  $\Phi$ -invariant Ricci property, we get that  $M$  has flat Ricci tensor.  $\square$

Finally, we established an application by identifying the necessary and sufficient condition for the locally conformal almost contact manifold to be  $\eta$ -Einstein manifold

**Theorem 7.** *Suppose that  $M$  is  $\mathcal{LCA}\mathcal{C}_S$ -manifold of class  $CT_1$  with  $\Phi$ -invariant Ricci tensor. Then the necessary and sufficient condition for  $M$  to be  $\eta$ -Einstein manifold is  $\sigma_d^a = E \delta_d^a$ , where  $E = \frac{2(1-n)\sigma_0^2}{n-2} - \frac{\alpha}{2n-1}$ .*

**P r o o f.** Suppose that  $M$  is  $\mathcal{LCA}\mathcal{C}_S$ -manifold of class  $CT_1$ . According to the Definition 11 and Lemma 5, we have

$$2(2\delta_{[c}^b \sigma_{d]}^a] - \delta_{[c}^a \delta_{d]}^b \sigma_0^2) - \frac{4}{2n-1} (r_{[d}^a \delta_{c]}^b) = 0. \quad (18)$$

Contracting (18) by indices  $(b, c)$ , we get

$$(n-2)\sigma_d^a + \delta_d^a \sigma_b^b - 2(1-n)\sigma_0^2 \delta_d^a - \frac{1}{2n-1} ((n-2)r_d^a + \delta_d^a r_b^b) = 0.$$

Then, we deduce

$$\sigma_d^a = \frac{2(1-n)\sigma_0^2\delta_d^a}{n-2} - \frac{r_d^a}{2n-1}. \quad (19)$$

Let  $M$  be  $\eta$ -Einstein manifold. We use the Definition 9, so the equation (19) becomes

$$\sigma_d^a = E\delta_d^a. \quad (20)$$

Conversely, from the equations (19) and (20), we have

$$r_d^a = \alpha\delta_d^a$$

where  $\alpha$  is the cosmological constant. Also

$$r_{00} = \alpha + \beta$$

where  $\alpha = -2n(\sigma_{00} + \sigma_0^2) - 2B_{hc}B^{ch} - 2(\sigma_c^c + \sigma^c\sigma_c) + 4\sigma^{[c}\delta_c^h]\sigma_h - \alpha$

According to  $\Phi$ -invariant of Ricci tensor, we obtain that  $M$  is  $\eta$ -Einstein manifold.  $\square$

#### REFERENCES

1. Abood H. M., Al-Hussaini F. H. Locally conformal almost cosymplectic manifold of  $\Phi$ -holomorphic sectional conharmonic curvature tensor, *European Journal of Pure and Applied Mathematics*, 2018, vol. 11, no. 3, pp. 671–681. <https://doi.org/10.29020/nybg.ejpam.v11i3.3261>
2. Abood H. M., Al-Hussaini F. H. Constant curvature of a locally conformal almost cosymplectic manifold, *AIP Conference Proceedings*, 2019, vol. 2086, issue 1, 030003. <https://doi.org/10.1063/1.5095088>
3. Abood H. M., Al-Hussaini F. H. On the conharmonic curvature tensor of a locally conformal almost cosymplectic manifold, *Communications of the Korean Mathematical Society*, 2020, vol. 35, no. 1, pp. 269–278. <https://doi.org/10.4134/CKMS.c190003>
4. Asghari N., Taleshian A. On the conharmonic curvature tensor of Kenmotsu manifolds, *Thai Journal of Mathematics*, 2014, vol. 12, no. 3, pp. 525–536. <http://thaijmath.in.cmu.ac.th/index.php/thaijmath/article/view/410>
5. Blair D. E. The theory of quasi-Sasakian structures, *Journal of Differential Geometry*, 1967, vol. 1, no. 3–4, pp. 331–345. <https://doi.org/10.4310/jdg/1214428097>
6. Blair D. E. *Riemannian geometry of contact and symplectic manifolds*, Boston, MA: Birkhäuser, 2010. <https://doi.org/10.1007/978-0-8176-4959-3>
7. Cartan E. *Riemannian geometry in an orthogonal frame*, Singapore: World Scientific, 2001.
8. Chanyal S. K., Upreti J. Conharmonic curvature tensor on  $(\kappa, \mu)$ -contact metric manifold, *An. Ştiinţ. Univ. Al. I. Cuza Iaşi Mat. (N.S.)*, 2016, vol. 2, issue F2, pp. 681–694. [https://www.math.uaic.ro/~annalsmath/new/?page\\_id=351](https://www.math.uaic.ro/~annalsmath/new/?page_id=351)
9. Chinea D., Marrero J. C. Classification of almost contact metric structures, *Revue Roumaine de Mathématiques Pures et Appliquées*, 1992, vol. 37, no. 3, pp. 199–211.
10. Dwivedi M. K., Kim J.-S. On conharmonic curvature tensor in  $K$ -contact and Sasakian manifolds, *Bulletin of the Malaysian Mathematical Sciences Society. Second Series*, 2011, vol. 34, no. 1, pp. 171–180.
11. Ghosh S., De U. C., Taleshian A. Conharmonic curvature tensor on  $N(K)$ -contact metric manifolds, *ISRN Geometry*, 2011, vol. 2011, article ID 423798, 11 p. <https://doi.org/10.5402/2011/423798>
12. Goldberg S. I., Yano K. Integrability of almost cosymplectic structures, *Pacific Journal of Mathematics*, 1969, vol. 31, no. 2, pp. 373–382. <https://doi.org/10.2140/pjm.1969.31.373>
13. Ishii Y. On conharmonic transformation, *Tensor, New Ser.*, 1957, vol. 7, pp. 73–80. <https://zbmath.org/?q=an:0079.15702>

14. Kharitonova S. V. On the geometry of locally conformal almost cosymplectic manifolds, *Mathematical Notes*, 2009, vol. 86, no. 1, pp. 121–131. <https://doi.org/10.1134/S0001434609070116>
15. Kirichenko V. F. *Differentsial'no-geometricheskie struktury na mnogoobraziyakh* (Differential-geometric structures on manifolds), Odessa: Pechatnyi Dom, 2013.
16. Kirichenko V. F., Rustanov A. R. Differential geometry of quasi Sasakian manifolds, *Sbornik: Mathematics*, 2002, vol. 193, no. 8, pp. 1173–1201. <https://doi.org/10.1070/SM2002v193n08ABEH000675>
17. Olszak Z. Locally conformal almost cosymplectic manifolds, *Colloquium Mathematicum*, 1989, vol. 57, no. 1, pp. 73–87. <https://doi.org/10.4064/cm-57-1-73-87>
18. Taleshian A., Prakasha D. G., Vikas K., Asghari N. On the conharmonic curvature tensor of  $LP$ -Sasakian manifolds, *Palestine Journal of Mathematics*, 2016, vol. 5, no. 1, pp. 177–184.
19. Volkova E. S. Curvature identities of normal manifolds of Killing type, *Mathematical Notes*, 1997, vol. 62, no. 3, pp. 296–305. <https://doi.org/10.1007/BF02360870>

Received 12.02.2020

Al-Hussaini Farah Hassan, University of Basrah, Basrah, Iraq.

E-mail: [farahalhussaini14@yahoo.com](mailto:farahalhussaini14@yahoo.com)

Abood Habeeb Mtashar, PhD, Professor, University of Basrah, Basrah, Iraq.

E-mail: [iraqsafwan2006@gmail.com](mailto:iraqsafwan2006@gmail.com)

**Citation:** F. H. Al-Hussaini, H. M. Abood. Quasi invariant conharmonic tensor of special classes of locally conformal almost cosymplectic manifold, *Vestnik Udmurtskogo Universiteta. Matematika. Mekhanika. Komp'yuternye Nauki*, 2020, vol. 30, issue 2, pp. 147–157.

**Ф. Х. Аль-Хусейни, Х. М. Абуд**

**Квазиинвариантный конгармонический тензор специальных классов локально конформного почти косимплектического многообразия**

*Ключевые слова:* локально конформное почти косимплектическое многообразие, тензор конгармонической кривизны,  $\Phi$ -квазиинвариант, конгармонический  $\Phi$ -параcontact.

УДК 514.7

DOI: [10.35634/vm200201](https://doi.org/10.35634/vm200201)

В работе описывается классификация локально конформного почти косимплектического многообразия ( $\mathcal{LCA}_S$ -многообразия) в соответствии с тензором конгармонической кривизны. В частности, были получены необходимые условия  $\Phi$  инвариантности тензора конгармонической кривизны на  $\mathcal{LCA}_S$ -многообразии классов  $CT_i$ ,  $i = 1, 2, 3$ . Кроме того, доказано, что любое  $\mathcal{LCA}_S$ -многообразие класса  $CT_1$  оказывается конгармоническим и  $\Phi$ -параcontactным.

Поступила в редакцию 12.02.2020

Аль-Хусейни Фара Хасан, университет Басры, Ирак, г. Басра.

E-mail: [farahalhussaini14@yahoo.com](mailto:farahalhussaini14@yahoo.com)

Абуд Хабиб Мташар, PhD, профессор, университет Басры, Ирак, г. Басра.

E-mail: [iraqsafwan2006@gmail.com](mailto:iraqsafwan2006@gmail.com)

**Цитирование:** Ф. Х. Аль-Хусейни, Х. М. Абуд. Квазиинвариантный конгармонический тензор специальных классов локально конформного почти косимплектического многообразия // Вестник Удмуртского университета. Математика. Механика. Компьютерные науки. 2020. Т. 30. Вып. 2. С. 147–157.